

Novel biology models in gut physiology in dairy cattle

B.M. Goetz, S. Rodriguez-Jimenez, M. A. Abeyta, J. Opgenorth, E. J. Mayorga,
and L. H. Baumgard
Department of Animal Science
Iowa State University

Introduction

Suboptimal milk yield limits the U.S. dairy industry's productive competitiveness, marginalizes efforts to reduce inputs into food production, and increases animal agriculture's carbon footprint. There are a variety of circumstances in a cow's life which result in hindered productivity including heat stress, ketosis, rumen and hindgut acidosis, feed restriction, and psychological stress associated with normal animal practices (i.e., pen changes, weaning, shipping). Although these insults have different origins, a commonality among them is increased production of inflammatory biomarkers and markedly altered nutrient partitioning. We and others have generated convincing data strongly implicating intestinally derived lipopolysaccharide (LPS) as sometimes being the culprit in these situations.

Heat Stress

During heat stress (HS), blood flow is diverted from the viscera to the periphery in an attempt to dissipate heat, and this leads to intestinal hypoxia (Hall et al., 1999). Enterocytes are particularly sensitive to hypoxia and nutrient restriction (Rollwagen et al., 2006), resulting in ATP depletion and increased oxidative and nitrosative stress (Hall et al., 2001). This contributes to tight junction dysfunction and gross morphological changes that ultimately reduce intestinal barrier function (Lambert et al., 2002; Pearce et al., 2013), resulting in increased passage of luminal content into portal and systemic blood (Hall et al., 2001; Pearce et al., 2013). Endotoxin, otherwise referred to as LPS, is a glycolipid embedded in the outer membrane of Gram-negative bacteria, which is abundant and prolific in luminal content, and is a well-characterized potent immune stimulator in multiple species (Berczi et al., 1966; Giri et al., 1990; Tough et al., 1997). Immune system activation occurs when LPS binding protein (LBP) initially binds LPS and together with CD14 and TLR4 delivers LPS for removal and detoxification, thus LBP is frequently used as a biomarker for LPS infiltration (Ceciliani et al., 2012). For a detailed description of how livestock and other species detoxify LPS see our recent review (Mani et al., 2012). Endotoxin infiltration into the bloodstream during HS, which was first observed by Graber et al. (1971), is common among heat stroke patients (Leon, 2007) and is thought to play a central role in heat stroke pathophysiology as survival increases when intestinal bacterial load is reduced or when plasma LPS is neutralized (Bynum et al., 1979; Gathiram et al., 1987). It is remarkable how animals suffering from heat stroke or severe endotoxemia share many physiological and metabolic similarities to HS, such as an increase in circulating insulin (Lim et al., 2007). Intramammary LPS infusion increased (~2 fold) circulating insulin in lactating cows (Waldron et al., 2006). In addition, we intravenously infused LPS into growing calves and pigs and demonstrated >10 fold increase in circulating insulin (Rhoads et al., 2009; Kvidera et al., 2016, 2017c). Interestingly, increased insulin occurs prior to increased inflammation and the temporal pattern agrees with our previous *in vivo* data and a recent *in vitro* report (Bhat et al., 2014) suggesting LPS stimulates insulin secretion, either directly or via GLP-1 (Kahles et al., 2014). The possibility that LPS increases insulin secretion likely explains the hyperinsulinemia we have repeatedly reported in a variety of HS agriculture models (Baumgard and Rhoads, 2013). Again, the increase in insulin during both HS and immunoactivation is energetically difficult to explain as feed intake is severely depressed in both experiments.

Ketosis and the Transition Period

Recently, the concept that LPS impacts normal nutrient partitioning and potentially contributes to metabolic maladaptation to lactation has started to receive attention. Although LPS itself has not been the primary causative focus, general inflammation has been the topic of investigations. Increased inflammatory markers following parturition have been reported in cows (Ametaj et al., 2005; Bionaz et al., 2007; Bertoni et al., 2008; Humblet et al., 2006; Mullins et al., 2012). Presumably, the inflammatory state following calving disrupts normal nutrient partitioning and is detrimental to productivity (Bertoni et al., 2008), and this assumption was recently reinforced when TNF α infusion decreased productivity (albeit without overt changes in metabolism; Yuan et al., 2013; Martel et al., 2014). Additionally, in late-lactation cows, injecting TNF α increased (>100%) liver TAG content without a change in circulating NEFA (Bradford et al., 2009). Our recent data demonstrates increased inflammatory markers in cows diagnosed with ketosis only and no other health disorders (i.e. the inflammation was not apparently due to mastitis or metritis). In comparison with healthy controls, ketotic cows had increased circulating LPS prior to calving and post-partum acute phase proteins such as LBP, serum amyloid A, and haptoglobin were also increased (Figure 1; Abuajamieh et al., 2016a). However, even seemingly healthy cows experience some degree of inflammation postpartum (Humblet et al., 2006). The magnitude and persistency of the inflammatory response seems to be predictive of transition cow performance (Bertoni et al., 2008; Bradford et al., 2015; Trevisi and Minuti, 2018). Endotoxin can originate from a variety of locations, and obvious sources in transitioning dairy cows include the uterus (metritis) and mammary gland (mastitis) (Mani et al., 2012). Additionally, we believe intestinal hyperpermeability may also be responsible for periparturient inflammation in dairy cows as many of the characteristic responses (rumen acidosis, decreased feed intake, and psychological stress) occurring during this time can compromise gut barrier function.




Figure 1. Markers of inflammation in healthy (solid line) and ketotic (dashed line) transition cows.

As aforementioned, mild inflammation is observed even in cows which seemingly complete the transition period successfully, suggesting that some level of inflammation plays an important role in cow health. In fact, previous reports have demonstrated that blocking endogenous inflammation (via administration of non-steroidal anti-inflammatory drugs [NSAID]) can increase the incidence of negative health outcomes (i.e., fever, stillbirth, retained placenta, metritis) and reduce productivity (Schwartz et al., 2009; Newby et al., 2013, 2017). Beneficial effects of NSAIDs have been observed on production performance (Carpenter et al., 2016), but inconsistencies exist (Priest et al., 2013; Meier et al., 2014) including how NSAIDs seemingly work better in specific parities (Farney et al., 2013) and interfere with fiber digestion (Carpenter et al., 2016) and compromise feed intake (Carpenter et al., 2017). Although NSAIDs may be an effective prophylactic strategy during the periparturient period, further research is necessary to determine the timing of administration and type and dose of NSAID that is most effective at improving health. Alternatively, administering a chemokine (anti or even pro-inflammatory) may hold promise in improving transition cow performance.

Rumen and Hindgut Acidosis

A transitioning dairy cow undergoes a dietary shift from a high forage to a high concentrate ration post-calving. This has the potential to induce rumen acidosis (RA) as increases in fermentable carbohydrates and DMI stimulate the buildup of short chain fatty acids and lactic acid (Nocek, 1997; Enemark, 2008). Rumen acidosis has direct and ancillary consequences accompanied by various production issues (decreased DMI, reduced milk yield, milk fat depression) and health challenges such as laminitis, liver abscesses, and potentially death (Nocek, 1997; Kleen, 2003). The mechanisms linking RA and the development of health disorders are not entirely clear, however, recent literature has indicated that inflammation associated with epithelial damage and consequential LPS translocation are at least partially responsible for production losses associated with RA (Gozho, et al., 2005; Khafipour, et al., 2009). Although many hypothesize LPS translocation occurs at the rumen epithelium directly (Guo et al., 2017; Minuti et al., 2014), others point towards LPS translocation in the hindgut to be a potential source of peripheral inflammation (Li et al., 2012). Interestingly, when RA was induced using either alfalfa pellets or high-grain diets, increased peripheral inflammation was only observed in the high-grain group, irrespective of rumen acidotic conditions being similar between the two treatments (Khafipour et al., 2009a,b). It was hypothesized that the grain supplemented group likely had increased starch flow to the hindgut, and therefore, increased fermentation that could potentially lead to hindgut acidosis and LPS translocation across the large intestine. However, we were unable to recreate production losses and systemic inflammation when we abomasally infused 500 g/d of resistant starch (Piantoni et al., 2018) or even 4 kg/d of purified corn starch (Abeyta et al., 2019). Both of our aforementioned experiments were accompanied with marked reductions in fecal pH so it is unlikely that large intestinal acidosis per se is the specific reason for systemic inflammation described in the previous reports (Li et al., 2012, Khafipour et al., 2009a,b). Regardless, we recently reported that cows with the largest decrease in fecal pH post-calving consumed less feed, produced less milk, had a larger acute phase protein response and had increased NEFA and BHB compared to cows that had a mild decrease in fecal pH following parturition (Rodriguez-Jimenez et al., 2019). Clearly, our current understanding of how hind-gut acidosis impacts the immune system and ultimately periparturient productivity is woeful.

Feed Restriction and Psychological Stress

Stress associated with feed restriction along with several other regular production practices (e.g., heat stress, weaning, transportation, overcrowding, restraint, social isolation/mixing) is frequently encountered in animal agriculture (Chen et al., 2015) and is associated with gastrointestinal hyperpermeability. In fact, we have repeatedly reported reduced intestinal barrier integrity in pigs pair-fed to their HS counterparts (Pearce et al., 2013; Sanz-Fernandez et al., 2014). Furthermore, we recently demonstrated shortened ileum villous height and crypt depth (Kvidera et al., 2017d) as well as increased appearance of the intestinal permeability marker Cr-EDTA (Horst and Baumgard, unpublished), indicating reduced intestinal health in cows fed 40% of ad libitum intake. Recent literature indicates that the corticotropin releasing factor (CRF) system may be the mechanism involved in stress-induced leaky gut (Wallon et al., 2008; Vanuytsel et al., 2014). The CRF and other members of the CRF signaling family including urocortin (1, 2,

and 3) and their G-protein couple receptors CRF1 and CRF2, have been identified as the main mediators of the stress-induced intestinal changes including inflammation, altered intestinal motility and permeability, as well as shifts in ion, water, and mucus secretion and absorption (as reviewed by Rodiño-Janeiro et al., 2015). These alterations appear to be regulated in large part by intestinal mast cells (Santos et al., 2000). Mast cells are important mediators of both innate and adaptive immunity and express receptors for the neuropeptides CRF1 and CRF2, which may in part explain the association between emotional stress and intestinal dysfunction (Smith et al., 2010; Ayyadurai et al., 2017). Furthermore, mast cells synthesize a variety of pro-inflammatory mediators (i.e., IFN- γ and TNF- α) that are released upon activation, mainly via degranulation (de Punder and Pruijboom, 2015). Excessive mast cell degranulation plays an important role in the pathogenesis of different intestinal inflammatory disorders (Santos et al., 2000; Smith et al., 2010). A better understanding of the role psychosocial stress plays on the initiation of different intestinal disorders in livestock is of obvious interest for multiple animal agriculture systems.

Metabolism of Inflammation

LPS-induced inflammation has an energetic cost which redirects nutrients away from anabolic processes that support milk and muscle synthesis (see review by Johnson 1997, 1998) and thus compromises productivity. Upon activation, most immune cells become obligate glucose utilizers via a metabolic shift from oxidative phosphorylation to aerobic glycolysis (not anaerobic glycolysis typically learned about in biochemistry classes), a process known as the Warburg effect (Figure 2).

This metabolic shift allows for rapid ATP production and synthesis of important intermediates which support proliferation and production of reactive oxygen species (Calder et al., 2007; Pálsson-McDermott and O'Neill, 2013). In an effort to facilitate glucose uptake, immune cells become more insulin sensitive and increase expression of GLUT3 and GLUT4 transporters (Maratou et al., 2007; O'Boyle et al., 2012), whereas peripheral tissues become insulin resistant (Poggi et al., 2007; Liang et al., 2013). Furthermore, metabolic adjustments including hyperglycemia or hypoglycemia (depending upon the stage and severity of infection), increased circulating insulin and glucagon, skeletal muscle catabolism and subsequent nitrogen loss (Figure 3; Wannemacher et al., 1980), and hypertriglyceridemia (Filkins, 1978; Wannemacher et al., 1980; Lanza-Jacoby et al., 1998; McGuinness, 2005) occur. Interestingly, despite hypertriglyceridemia, circulating BHB often decreases following LPS administration (Waldron et al., 2003a,b; Graugnard et al., 2013; Kvidera et al., 2017a). The mechanism of LPS-induced decreases in BHB has not been fully elucidated, but may be explained by increased ketone oxidation by peripheral tissues (Zarrin et al., 2014). Collectively, these metabolic alterations are presumably employed to ensure adequate glucose delivery to activated leukocytes.




Figure 2. Metabolic pathway of a resting (A) vs. activated (B) leukocyte




Figure 3. LPS induced alterations in peripheral metabolism.

Energetic Cost of Immune Activation

The energetic costs of immunoactivation are substantial, but the ubiquitous nature of the immune system makes quantifying the energetic demand difficult. Our group recently employed a series of LPS-euglycemic clamps to quantify the energetic cost of an activated immune system. Using this model, we

estimated approximately 1 kg of glucose is used by an intensely activated immune system during a 12 hour period in lactating dairy cows. Interestingly, on a metabolic body weight basis the amount of glucose utilized by LPS-activated immune system in mid- and late-lactation cows, growing steers and growing pigs were 0.64, 1.0, 0.94, 1.0, and 1.1 g glucose/kg BW^{0.75}/h, respectively; Kvidera et al., 2016, 2017b,c, Horst et al., 2018, 2019). A limitation to our model is the inability to account for liver's contribution to the circulating glucose pool (i.e., glycogenolysis and gluconeogenesis). However, both glycogenolytic and gluconeogenic rates have been shown to be increased during infection (Spitzer et al., 1985; Waldron et al., 2003b) and Waldron et al. (2006) demonstrated that ~87 g of glucose appeared in circulation from these processes. Furthermore, we have observed both increased circulating glucagon and cortisol (stimulators of hepatic glucose output) following LPS administration (Horst et al., 2019) suggesting we are underestimating the energetic cost of immunoactivation. The reprioritization of glucose trafficking during immunoactivation has particular consequences during lactation as it requires ~72 g of glucose for synthesizing 1 kg milk (Kronfeld, 1982).

Increased immune system glucose utilization occurs simultaneously with infection-induced decreased feed intake: this coupling of enhanced nutrient requirements with hypophagia obviously decrease the amount of nutrients available for the synthesis of valuable products (milk, meat, fetus, wool, etc.). We and others have now demonstrated that HS, rumen acidosis, and psychological stress increase circulating markers of endotoxin and inflammation. We believe that the circulating LPS originates from the intestine (small or large) and initiates an immune response. This activated systemic immune response reprioritizes the hierarchy of glucose utilization and milk synthesis is consequently deemphasized.

Nutritional Mitigation Strategies: The Role of Zinc (Zn) Supplementation

Potential dietary mitigation strategies aimed at improving gut health are currently of great interest, especially considering the numerous stressors (i.e., heat stress, feed restriction, acidosis) that potentially impact intestinal permeability. Zinc is an essential nutrient which is crucial for maintaining epithelial integrity (i.e., mammary, uterine, intestinal) and regulating the renewal of damaged epithelium (Alam et al., 1994). Zinc was first demonstrated to improve intestinal "health" in human leaky gut models (Alam et al., 1994; Rodriguez et al., 1996; Sturniolo et al., 2001), and we extended this to improved metrics of intestinal permeability in a variety of farm animal stress models including heat stress (Sanz-Fernandez et al., 2014; Pearce et al., 2015; Abuajamieh et al., 2016b) and feed restriction (Horst and Baumgard, unpublished using Zn hydroxychloride). Additionally, we observed altered febrile, cytokine, and acute phase protein responses during heat stress (Sanz-Fernandez et al., 2014; Pearce et al., 2015; Abuajamieh et al., 2016b; Mayorga et al., 2018) and in response to LPS administration (Horst et al., 2019) with dietary Zn supplementation. Presumably the aforementioned changes in inflammatory variables are indicative of a blunted immune response (because of improved intestinal barrier function). Therefore, Zn as a dietary supplement appears to be a promising avenue to improve gut health and to ameliorate alimentary canal associated inflammation.

Conclusion

There are various situations in an animal's life that hinder production performance (i.e., heat stress, feed restriction, rumen acidosis, etc.) and we suggest, based upon the literature and on our supporting evidence, that LPS (of intestinal origin) may be the common culprit in these circumstances. Immune activation in response to LPS markedly alters nutrient partitioning as a means of fueling the immune response. More research is still needed to understand the mechanisms and consequences of intestinal permeability and associated inflammation in order to provide foundational information for developing strategies aimed at maintaining productivity.

*Parts of this manuscript were first published in the proceedings of the 2016, 2017 and 2018 Southwest Nutrition Conference in Tempe AZ and 2019 Cornell Nutrition Conference.

Literature Cited

- Abeyta, M. A., E. A. Horst, E. J. Mayorga, B. M. Goetz, M. Al-Qaisi, C. S. McCarthy, M. R. O'Neil, B. C. Dooley, P. Piantoni, G. F. Schroeder, H. A. Ramirez-Ramirez, and L. H. Baumgard. 2019. Effects of hindgut acidosis on metabolism, inflammation and production in dairy cows consuming a standard lactation diet. *J. Dairy Sci.* (Suppl. 1): 270.
- Abujamieh, M., S.K. Kvidera, M.V. Fernandez, A. Nayeri, N.C. Upah, E.A. Nolan, S.M. Lei, J.M. DeFrain, H.B. Green, K.M. Schoenberg, E.B. Trout, and L.H. Baumgard. 2016a. Inflammatory biomarkers are associated with ketosis in periparturient Holstein cows. *Res. Vet. Sci.* 109:81-85. doi:10.1016/j.rvsc.2016.09.015
- Abujamieh, M., S. K. Kvidera, E. A. Horst, E. J. Mayorga, J. T. Seibert, J. S. Johnson, J. W. Ross, M. A. Al-Qaisi, P. J. Gordon, J. DeFrain, R.P. Rhoads, and L. H. Baumgard. 2016b. The effects of zinc amino acid complex on biomarkers of gut integrity and metabolism in heat-stressed steers. *J. Dairy Sci.* 99(E-Suppl. 1):1175
- Alam, A. N., S. A. Sarker, and M. A. Wahed. 1994. Enteric protein loss and intestinal permeability changes in children during acute shigellosis and after recovery: effect of zinc supplementation. *Gut* 35:1707–1711
- Ametaj, B. N., B. J. Bradford, G. Bobe, R. A. Nafikov, Y. Lu, J. W. Young, and D. C. Beitz. 2005. Strong relationships between mediators of the acute phase response and fatty liver in dairy cows. *Can. J. Anim. Sci.* 85:165–175.
- Ayyadurai, S., A. J. Gibson, S. D'Costa, E. L. Overman, L. J. Sommerville, A. C. Poopal, E. Mackey, Y. Li, and A. J. Moeser. 2017. Corticotropin-releasing factor receptor subtype 1 is a critical modulator of mast cell degradation and stress-induced pathophysiology. *J. Leukoc. Biol.* 102:1299-1312.
- Baumgard, L. H. and R. P. Rhoads. 2013. Effects of heat stress on postabsorptive metabolism and energetics. *Annu. Rev. Anim. Biosci.* 1:311–337.
- Berczi, I., L. Bertok, and T. Bereznoi. 1966. Comparative studies on the toxicity of *Escherichia coli* lipopolysaccharide endotoxin in various animal species. *Can. J. of Microbiol.* 12:1070-1071.
- Bertoni, G., E. Trevisi, X. Han, and M. Bionaz. 2008. Effects of inflammatory conditions on liver activity in puerperium period and consequences for performance in dairy cows. *J. Dairy Sci.* 91:3300–3310.
- Bhat, U. G., V. Ilievski, T. G. Unterman, and K. Watanabe. 2014. *Porphyromonas gingivalis* lipopolysaccharide (LPS) upregulates insulin secretion from pancreatic beta cells line MIN6. *J. Periodontol.* 85:1629–1636.
- Bionaz, M., E. Trevisi, L. Calamari, F. Librandi, A. Ferrari, and G. Bertoni. 2007. Plasma paraoxonase, health, inflammatory conditions, and liver function in transition dairy cows. *J. Dairy Sci.* 90:1740-1750.
- Bradford, B. J., L. K. Mamedova, J. E. Minton, J. S. Drouillard, and B. J. Johnson. 2009. Daily injection of tumor necrosis factor- α increases hepatic triglycerides and alters transcript abundance of metabolic genes in lactating dairy cattle. *J. Nutr.* 139:1451–1456.
- Bradford, B.J., Yuan, K., Farney, J.K., Mamedova, L.K., Carpenter, A.J., 2015. Invited review: Inflammation during the transition to lactation: New adventures with an old flame. *J. Dairy Sci.* 98, 6631-6650.

- Bynum, G., J. Brown, D. Dubose, M. Marsili, I. Leav, T. G. Pistole, M. Hamlet, M. LeMaire, and B. Caleb. 1979. Increased survival in experimental dog heatstroke after reduction of gut flora. *Aviat. Space Environ. Med.* 50:816-819.
- Calder, P. C., G. Dimitriadis, and P. Newsholme. 2007. Glucose metabolism in lymphoid and inflammatory cells and tissues. *Curr. Opin. Clin. Nutr. Metab. Care.* 10:531-540.
- Carpenter, A. J., C. M. Ylloja, C. F. Vargas, L. K. Mamedova, L. G. Mendonça, J. F. Coetzee, L. C. Hollis, R. Gehring, and B. J. Bradford. 2016. Hot topic: Early postpartum treatment of commercial dairy cows with nonsteroidal anti-inflammatory drugs increases whole-lactation milk yield. *J. Dairy Sci.* 99:672-679.
- Carpenter, A.J., C.M. Ylloja, and B.J. Bradford. 2016. Early postpartum administration of sodium salicylate to multiparous dairy cattle is associated with alterations in feeding behavior up to 120 d in milk. *J. Anim Sci.* 94. Suppl. 5. 531 (abst).
- Carpenter, A.J., C.F.V. Rodriguez, J.A.B. Jantz, and B.J. Bradford. 2017. Sodium salicylate negatively affects rumen fermentation in vitro and in situ. *J. Dairy Sci.* 100:1935-1939.
- Ceciliani, F., J.J. Ceron, P.D. Eckersall, and H. Sauerwein. 2012. Acute phase proteins in ruminants. *J. Proteomics.* 75:4207-4231.
- Chen, Y., R. Arsenault, S. Napper, and P. Griebel. 2015. Models and methods to investigate acute stress responses in cattle. *Animals (Basel).* 5:1268-1295.
- de Punder, K., and L. Pruimboom. 2015. Stress induces endotoxemia and low-grade inflammation by increasing barrier permeability. *Front. Immunol.* 6:223.
- Enemark, J. M. D. 2008. The monitoring, prevention and treatment of sub-acute ruminal acidosis (SARA): a review. *Vet. J.* 176:32-43.
- Farney, J. K., L. K. Mamedova, J. F. Coetzee, J. E. Minton, L. C. Hollis, and B. J. Bradford. 2013. Sodium salicylate treatment in early lactation increases whole-lactation milk and milk fat yield in mature dairy cows. *J. Dairy Sci.* 96:7709-7718.
- Filkins, J. P. 1978. Phases of glucose dyshomeostasis in endotoxemia. *Circ. Shock* 5:347-355.
- Gathiram, P., M. T. Wells, J. G. Brock-Utne, and S. L. Gaffin. 1987. Antilipoplysaccharide improves survival in primates subjected to heat stroke. *Circ. Shock* 23:157-164.
- Giri, S. N., P. Emau, J. S. Cullor, G. H. Stabenfeldt, M. L. Bruss, R. H. Bondurant, and B. I. Osburn. 1990. Effects of endotoxin infusion on circulating levels of eicosanoids, progesterone, cortisol, glucose and lactic acid, and abortion in pregnant cows. *Vet. Microbiol.* 21:211-231.
- Gozho, G. N., J. C. Plaizier, D. O. Krause, A. D. Kennedy, and K. M. Wittenberg. 2005. Subacute Ruminal Acidosis Induces Ruminal Lipopolysaccharide Endotoxin Release and Triggers an Inflammatory Response. *J. Dairy Sci.* 88:1399-1403
- Graber, C. D., R. B. Reinhold, J.G. Breman, R. A. Harley, and G. R. Hennigar. 1971. Fatal heat stroke. Circulating endotoxin and gram-negative sepsis as complications. *JAMA.* 216:1195-1196.
- Graugnard, D. E., K. M. Moyes, E. Trevisi, M. J. Khan, D. Keisler, J. K. Drackley, G. Bertoni, and J. J. Looor. 2013. Liver lipid content and inflammometabolic indices in periparturient dairy cows are altered in response to preparturient energy intake and postparturient intramammary inflammatory challenge. *J. Dairy Sci.* 96:918-935.

- Guo, J., G. Chang, K. Zhang, L. Xu, D. Jin, M. S. Bilal, and X. Shen. 2017. Rumen-derived lipopolysaccharide provoked inflammatory injury in the liver of dairy cows fed a high-concentrate diet. *Oncotarget*. 8(29):46769-46780.
- Hall, D. M., K. R. Baumgardner, T. D. Oberley, and C. V. Gisolfi. 1999. Splanchnic tissues undergo hypoxic stress during whole body hyperthermia. *Am. J. Physiol.* 276:G1195-G1203.
- Hall, D. M., G. R. Buettner, L. W. Oberley, L. Xu, R. D. Matthes, and C. V. Gisolfi. 2001. Mechanisms of circulatory and intestinal barrier dysfunction during whole body hyperthermia. *Am. J. Physiol. Heart Circ. Physiol.* 280:H509– H521.
- Horst, E. A., S. K. Kvidera, E. J. Mayorga, C. S. Shouse, M. Al-Qaisi, M. J. Dickson, J. Ydstie, H. A. Ramirez Ramirez, A. F. Keating, D. J. Dickson, K. E. Griswold, and L. H. Baumgard. 2018a. Effect of chromium on bioenergetics and leukocyte dynamics following immunoactivation in lactating Holstein cows. *J. Dairy Sci.* 101:5515-5530.
- Horst, E. A., E. J. Mayorga, S. L. Portner, M. Al-Qaisi, C. S. McCarthy, M. A. Abeyta, B. M. Goetz, H. A. Ramirez-Ramirez, D. H. Kleinschmit, and L. H. Baumgard. 2019. Effects of dietary zinc on energetic requirements of an activated immune system following lipopolysaccharide challenge in lactating cows. *J. Dairy Sci.* (accepted).
- Humblet, M. F., H. Guyot, B. Boudry, F. Mbayahi, C. Hanzen, F. Rollin, and J. M. Godeau. 2006. Relationship between haptoglobin, serum amyloid A, and clinical status in a survey of dairy herds during a 6-month period. *Vet. Clin. Pathol.* 35:188–193.
- Johnson, R. W. 1997. Inhibition of growth by pro-inflammatory cytokines: an integrated view. *J Anim. Sci.* 75: 1244-1255.
- Johnson, R. W. 1998. Immune and endocrine regulation of food intake in sick animals. *Domest. Animal Endo.* 15: 309-319.
- Kahles, F., C. Meyer, J. Möllmann, S. Diebold, H.M. Findeisen, C. Leberherz, C. Trautwein, A. Koch, F. Tacke, N. Marx, and M. Lehrke. 2014. GLP-1 Secretion Is Increased by Inflammatory Stimuli in an IL-6–Dependent Manner, Leading to Hyperinsulinemia and Blood Glucose Lowering. *Diabetes.* 63:3221-3229.
- Khafipour, E., D.O. Krause, and J.C. Plaizier. 2009a. A grain-based subacute ruminal acidosis challenge causes translocation of lipopolysaccharide and triggers inflammation. *J. Dairy Sci.* 92:1060-1070.
- Khafipour, E., D. O. Krause, and J. C. Plaizier. 2009b. Alfalfa pellet-induced subacute ruminal acidosis in dairy cows increases bacterial endotoxin in the rumen without causing inflammation. *J. Dairy Sci.* 92:1712-1724.
- Kleen, J. L., G. A. Hooijer, J. Rehage, and J. P. T. M. Noordhuizen. 2003. Subacute ruminal acidosis (SARA): a review. *J. Vet. Med.* 50:406-414.
- Kronfeld, D. S. 1982. Major metabolic determinants of milk volume, mammary efficiency, and spontaneous ketosis in dairy cows. *J. Dairy Sci.* 65:2204-2212.
- Kvidera, S. K., E. A. Horst, M. Abuajamieh, E. J. Mayorga, M. V. Sanz-Fernandez, and L. H. Baumgard. 2016. Technical note: A procedure to estimate glucose requirements of an activated immune system in steers. *J. Anim. Sci.* 94:4591-4599.
- Kvidera, S. K., M. J. Dickson, M. Abuajamieh, D. B. Snider, M. V. Sanz-Fernandez, J. S. Johnson, A. F. Keating, P. J. Gorden, H. B. Green, K. M. Schoenberg, and L.H. Baumgard. 2017a. Intentionally induced intestinal barrier dysfunction causes inflammation, affects metabolism, and reduces productivity in lactating Holstein cows. *J. Dairy Sci.* 100:4113-4127.

- Kvidera, S. K., E. A. Horst, M. Abuajamieh, E. J. Mayorga, M. V. Sanz-Fernandez, and L. H. Baumgard. 2017b. Glucose requirements of an activated immune system in lactating Holstein cows. *J. Dairy Sci.* 100:2360-2374.
- Kvidera, S. K., E. A. Horst, E. J. Mayorga, M. V. Sanz-Fernandez, M. Abuajamieh, and L. H. Baumgard. 2017c. Estimating glucose requirements of an activated immune system in growing pigs. *J. Anim. Sci.* 95:5020-5029.
- Kvidera, S. K., E. A. Horst, M. V. Sanz-Fernandez, M. Abuajamieh, S. Ganesan, P. J. Gorden, H. B. Green, K. M. Schoenberg, W. E. Trout, A. F. Keating, and L. H. Baumgard. 2017d. Characterizing effects of feed restriction and glucagon-like peptide 2 administration on biomarkers of inflammation and intestinal morphology. *J. Dairy Sci.* 100:9402-9417.
- Lambert, G. P., C. V. Gisolfi, D. J. Berg, P. L. Moseley, L. W. Oberley, and K. C. Kregel. 2002. Selected contribution: Hyperthermia-induced intestinal permeability and the role of oxidative and nitrosative stress. *J. Appl. Physiol.* 92:1750-1761.
- Lanza-Jacoby, S., H. Phetteplace, N. Sedkova, and G. Knee. 1998. Sequential alterations in tissue lipoprotein lipase, triglyceride secretion rates, and serum tumor necrosis factor alpha during *Escherichia coli* bacteremic sepsis in relation to the development of hypertriglyceridemia. *Shock* 9:46-51.
- Leon, L. R. 2007. Heat stroke and cytokines. *Prog. Brain Res.* 162:481-524.
- Li, S., E. Khafipour, D. O. Krause, A. Kroeker, J. C. Rodriguez-Lecompte, G. N. Gozho, and J. C. Plaizier. 2012. Effects of subacute ruminal acidosis challenges on fermentation and endotoxins in the rumen and hindgut of dairy cows. *J. Dairy Sci.* 95:294-303.
- Liang, H., S. E. Hussey, A. Sanchez-Avila, P. Tantiwong, and N. Musi. 2013. Effect of lipopolysaccharide on inflammation and insulin action in human muscle. *PLoS One* 8:e63983.
- Lim, C. L., G. Wilson, L. Brown, J. S. Coombes, and L. T. Mackinnon. 2007. Pre-existing inflammatory state compromises heat tolerance in rats exposed to heat stress. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 292:R186-194.
- Mani, V., T. E. Weber, L. H. Baumgard and N. K. Gabler. 2012. Growth and development symposium: endotoxin, inflammation, and intestinal function in livestock. *J. Anim. Sci.* 90:1452-1465.
- Maratou, E., G. Dimitriadis, A. Kollias, E. Boutati, V. Lambadiari, P. Mitrou, and S. A. Raptis. 2007. Glucose transporter expression on the plasma membrane of resting and activated white blood cells. *Eur. J. Clin. Invest.* 37:282-290.
- Mayorga, E. J., S. K. Kvidera, E. A. Horst, M. A. Al-Qaisi, M. J. Dickson, J. T. Seibert, S. Lei, Z. J. Rambo, M. E. Wilson, and L. H. Baumgard. 2018. Effects of zinc amino acid complex on biomarkers of gut integrity and metabolism during and following heat stress or feed restriction in pigs. *J. Anim. Sci.* 96:4173-4185.
- Martel, C. A., L. K. Mamedova, J. E. Minton, M. L. Jones, J. A. Carroll, and B. J. Bradford. 2014. Continuous low-dose infusion of tumor necrosis factor alpha in adipose tissue elevates adipose tissue interleukin 10 abundance and fails to alter metabolism in lactating dairy cows. *J. Dairy Sci.* 97:4897-4906.

- McGuinness, O. P. 2005. Defective glucose homeostasis during infection. *Annu. Rev. Nutr.* 25:9-35.
- Meier, S., N. V. Priest, C. R. Burke, J. K. Kay, S. McDougall, M. D. Mitchell, C. G. Walker, A. Heiser, J. J. Looor, and J. R. Roche. 2014. Treatment with a nonsteroidal antiinflammatory drug after calving did not improve milk production, health, or reproduction parameters in pasture-grazed dairy cows. *J. Dairy Sci.* 97:2932-2943.
- Minuti, A., S. Ahmed, E. Trevisi, F. Piccioli-Cappelli, G. Bertoni, N. Jahan, and P. Bani. 2014. Experimental acute rumen acidosis in sheep: Consequences on clinical, rumen, and gastrointestinal permeability conditions and blood chemistry. *J. Anim. Sci.* 92:3966-3977.
- Mullins, C. R., L. K. Mamedova, M. J. Brouk, C. E. Moore, H. B. Green, K. L. Perfield, J. F. Smith, J. P. Harner, and B. J. Bradford. 2012. Effects of monensin on metabolic parameters, feeding behavior, and productivity of transition dairy cows. *J. Dairy Sci.* 95:1323–1336.
- Newby, N.C., D.L. Pearl, S.J. Leblanc, K.E. Leslie, M.A. von Keyserlingk, and T.F. Duffield. 2013. Effects of meloxicam on milk production, behavior, and feed intake in dairy cows following assisted calving. *J. Dairy Sci.* 96:3682-3688.
- Newby, N.C., K. E. Leslie, H. D. Putnam Dingwell, D. F. Kelton, D. M. Weary, L. Neuder, S. T. Millman, and T. F. Duffield. 2017. The effects of periparturient administration of flunixin meglumine on the health and production of dairy cattle. *J. Dairy Sci.* 100:582-587.
- Nocek, J. E. 1997. Bovine acidosis: Implications on laminitis. *J. Dairy Sci.* 80:1005-1028.
- O'Boyle, N. J., G. A. Contreras, S. A. Mattmiller, and L. M. Sordillo. 2012. Changes in glucose transporter expression in monocytes of periparturient dairy cows. *J. Dairy Sci.* 95:5709-5719.
- Palsson-McDermott, E. M. and L. A. O'Neill. 2013. The Warburg effect then and now: from cancer to inflammatory diseases. *Bioessays* 35:965-973.
- Pearce, S. C., V. Mani, T. E. Weber, R. P. Rhoads, J. F. Patience, L. H. Baumgard, and N. K. Gabler. 2013. Heat stress and reduced plane of nutrition decreases intestinal integrity and function in pigs. *J. Anim. Sci.* 91:5183-5193.
- Pearce S. C., M. V. Sanz-Fernandez, J. Torrison, M. E. Wilson, L. H. Baumgard, and N. K. Gabler. 2015. Dietary zinc attenuates heat-stress induced changes in intestinal integrity and metabolism. *J. Anim. Sci.* 93:4702-4713.
- Piantoni, P., M.A. Abeyta, G.F. Schroeder, H.A. Ramirez-Ramirez, H.A. Tucker and L.H. Baumgard. 2018. Induction of leaky gut through feed restriction or abomasal infusion of resistant starch in healthy post-peak lactating cows. *J. Dairy Sci.* (Suppl. 2): 228.
- Poggi, M., D. Bastelica, P. Gual, M. A. Iglesias, T. Gremeaux, C. Knauf, F. Peiretti, M. Verdier, I. Juhán-Vague, J. F. Tanti, R. Burcelin, and M. C. Alessi. 2007. C3H/HeJ mice carrying a toll-like receptor 4 mutation are protected against the development of insulin resistance in white adipose tissue in response to a high-fat diet. *Diabetologia* 50:1267-1276.
- Priest, N. V., S. McDougall, C. R. Burke, J. R. Roche, M. Mitchell, K. L. McLeod, S. L. Greenwood, and S. Meier. 2013. The responsiveness of subclinical endometritis to a nonsteroidal anti-inflammatory drug in pasture-grazed dairy cows. *J. Dairy Sci.* 96:4323-4332.
- Rhoads, M. L., R. P. Rhoads, M. J. VanBaale, R. J. Collier, S. R. Sanders, W. J. Weber, B. A. Crooker, and L. H. Baumgard. 2009. Effects of heat stress and plane of nutrition on lactating Holstein cows: I. Production, metabolism, and aspects of circulating somatotropin. *J. Dairy Sci.* 92:1986-1997.

- Rodiño-Janeiro, B. K., C. Alonso-Cotner, M. Pigrau, B. Lobo, M. Vicario, and J. Santos. 2015. Role of corticotropin-releasing factor in gastrointestinal permeability. *J. Neurogastroenterol. Motil.* 21:33-50.
- Rodriguez-Jimenez, S., C. S. McCarthy, E. A. Horst, E. J. Mayorga, M. Al-Qaisi, M. A. Abeyta, B. M. Goetz, H. A. Ramirez-Ramirez, and L. H. Baumgard. 2019. Relationships between fecal pH and milk production, metabolism, and acute phase protein response in periparturient dairy cows. *J. Dairy Sci.* 102 (Suppl. 1): 402.
- Rodriguez, P., N. Darmon, P. Chappuis, C. Candalh, M. A. Blaton, C. Bouchaud, and M. Heyman. 1996. Intestinal paracellular permeability during malnutrition in guinea pigs: effect of high dietary zinc. *Gut* 39:416-422.
- Rollwagen, F. M., S. Madhavan, A. Singh, Y. Y. Li, K. Wolcott, and R. Maheshwari. 2006. IL-6 protects enterocytes from hypoxia-induced apoptosis by induction of bcl-2 mRNA and reduction of fas mRNA. *Biochem. Biophys. Res. Commun.* 347:1094-1098.
- Santos, J., M. Benjamin, P. C. Yang, T. Prior, and M. H. Perdue. Chronic stress impairs rat growth and jejunal epithelial barrier function: role of mast cells. 2000. *Am. J. Physiol. Gastrointest. Liver Physiol.* 278:G847-G854.
- Sanz-Fernandez, M.V., S. C. Pearce, N. K. Gabler, J. F. Patience, M. E. Wilson, M. T. Socha, J. L. Torrison, R. P. Rhoads, and L. H. Baumgard. 2014. Effects of supplemental zinc amino acid complex on gut integrity in heat-stressed growing pigs. *Animal.* 8:43-50.
- Schwartz, G., K. L. Hill, M. J. VanBaale, and L. H. Baumgard. 2009. Effects of flunixin meglumine on pyrexia and bioenergetics variables in postparturient dairy cows. *J. Dairy Sci.* 92:1963-1970.
- Smith, F., J. E. Clark, B. L. Overman, C. C. Tozel, J. H. Huang, J. E. F. Rivier, A. T. Blikslager, and A. J. Moeser. 2010. Early weaning stress impairs development of mucosal barrier function in the porcine intestine. *Am. J. Physiol. Gastrointest. Liver Physiol.* 298: G352-G363.
- Spitzer, J. A., K. M. Nelson, and R. E. Fish. 1985. Time course of changes in gluconeogenesis from various precursors in chronically endotoxemic rats. *Metabolism* 34:842-849.
- Sturniolo, G. C., V. Di Leo, A. Ferronato, A. D'Odorico, and R. D'Inca. 2001. Zinc supplementation tightens "leaky gut" in crohn's disease. *Inflamm. Bowel Dis.* 7:94-98.
- Tough, D. F., S. Sun, and J. Sprent. 1997. T cell stimulation in vivo by lipopolysaccharide (LPS). *J. Exp. Med.* 185:2089-2094.
- Trevisi, E., and A. Minuti. 2018. Assessment of the innate immune response in the periparturient cow. *Res. Vet. Sci.* 116:47-54.
- Vanuytsel, T., S. van Wanrooy, H. Vanheel, C. Vanormelingen, S. Verschueren, E. Houben, S. Salim Rasoel, J. Tóth, L. Holvoet, R. Farré, L. Van Oudenhove, G. Boeckxstaens, K. Verbeke, and J. Tack. 2014. Psychological stress and corticotropin-releasing hormone increase intestinal permeability in humans by a mast cell-dependent mechanism. *Gut* 63:1293-1299.
- Waldron, M. R., B. J. Nonnecke, T. Nishida, R. L. Horst, and T. R. Overton. 2003a. Effect of lipopolysaccharide infusion on serum macromineral and vitamin D concentrations in dairy cows. *J. Dairy Sci.* 86:3440-3446.
- Waldron, M. R., T. Nishida, B. J. Nonnecke, and T. R. Overton. 2003b. Effect of lipopolysaccharide on indices of peripheral and hepatic metabolism in lactating cows. *J. Dairy Sci.* 86:3447-3459.

- Waldron, M. R., A. E. Kulick, A. W. Bell, and T. R. Overton. 2006. Acute experimental mastitis is not causal toward the development of energy-related metabolic disorders in early postpartum dairy cows. *J. Dairy Sci.* 89:596-610.
- Wallon, C., P. C. Yang, A. V. Keita, A. C. Ericson, D. M. McKay, P. M. Sherman, M. H. Perdue, and J. D. Söderholm. 2008. Corticotropin-releasing hormone (CRH) regulates macromolecular permeability via mast cells in normal human colonic biopsies in vitro. *Gut* 57:50-58.
- Wannemacher, R. W., F. A. Beall, P. G. Canonico, R. E. Dinterman, C. L. Hadick, and H. A. Neufeld. 1980. Glucose and alanine metabolism during bacterial infections in rats and rhesus monkeys. *Metabolism* 29:201-212.
- Yuan, K., J. K. Farney, L. K. Mamedova, L. M. Sordillo, and B. J. Bradford. 2013. TNF α Altered Inflammatory Responses, Impaired Health and Productivity, but Did Not Affect Glucose or Lipid Metabolism in Early-Lactation Dairy Cows. *PloS One*. e80316.
- Zarrin, M., O. Wellnitz, H. A. van Dorland, J. J. Gross, and R. M. Bruckmaier. 2014. Hyperketonemia during lipopolysaccharide-induced mastitis affects systemic and local intramammary metabolism in dairy cows. *J. Dairy Sci.* 97:3531-3541.